

Prevalence of sexual activity and associated factors in men aged 75 to 95 years a cohort study

Author/s: Hyde, Z. Flicker, L. Hankey, G. J. Almeida, O. P. McCaul, K. A. Chubb, S. A. P. Yeap, B. B. | **Year:** 2010 | **Publication type:** Journal article | **Peer reviewed:** | **Topic area/s:** General Wellbeing

Reference: Hyde, Z., L. Flicker, G. J. Hankey, O. P. Almeida, K. A. McCaul, S. A. P. Chubb and B. B. Yeap (2010). "Prevalence of sexual activity and associated factors in men aged 75 to 95 years a cohort study." *Annals of Internal Medicine* 153(11): 693-702.

Key Words:

sex, men, health, wellbeing

Research aim:

To determine the proportion of older men who are sexually active and to explore factors predictive of sexual activity.

Results/Conclusion:

Sex was considered at least somewhat important by 48.8% (95% CI, 47.0% to 50.6%), and 30.8% (CI, 29.1% to 32.5%) had had at least 1 sexual encounter in the past 12 months. Living with a partner and having a non-English-speaking background were associated with increased odds of sexual activity.

Implications:

Cultural Group(s):

Not specific to CALD

Location of study:

Western Australia (Perth)

Age group:

75-95

Number included in study:

2,783

Type of participants:

Men aged 75-85

Research approach:

Quantitative

Type of data:

Primary

Secondary data sources used:

Specific scales or analytical techniques used:

Implications/ Recommendations:

Notes: